

# Associations of husbandry management factors with the new infection risk of bovine intramammary infections in lactation of dairy herds in Northern Germany

Matthias Gösling<sup>1</sup>, Doris Klocke<sup>1</sup>, Friederike Reinecke<sup>2</sup>, Veit Zoche-Golob<sup>3</sup>, Martin tho Seeth<sup>4</sup>, Jan-Hendrik Paduch<sup>1</sup>, Volker Krömker<sup>5</sup>

<sup>1</sup> Hannover University of Applied Sciences and Arts, Faculty II, Microbiology, Heisterbergallee 10a, 30453, Hannover, Germany

<sup>2</sup> Regional Council of Giessen, Dept. 51.2, 35531 Wetzlar, Germany

<sup>3</sup> German Federal Institute for Risk Assessment, Unit Epidemiology, Zoonoses and Antimicrobial Resistance, Diederdsdorfer Weg 1, 12277 Berlin, Germany

<sup>4</sup> Chamber of Agriculture Lower Saxony, Udder Health Service, Hermann-Ehlers-Straße 15, 26160 Bad Zwischenahn, Germany

<sup>5</sup> Department of Veterinary and Animal Sciences, University of Copenhagen, 1870 Frederiksberg C, Denmark

Date submitted: 06/03/2021

Date accepted: 24/03/2021

Volume/Page(s): 1-8

## Abstract

The purpose of this investigation was to compare herd-level new infection risk of bovine intramammary infections regarding husbandry management factors. The new infection risk was derived from cow-level somatic cell counts that were provided by monthly dairy herd improvement tests and collected over a period of 3 years and 4 months from 60 commercial dairy farms located in Lower Saxony, Germany. Factors of the management of the farm, the livestock and milking were generated as potential predicting variables. Based on the results of linear mixed models, the herd-level new infection risk was significantly associated with the pre-milking routine and the housing of fresh-lactating cows. When forestripping of every cow was included in the pre-milking routine, this had a beneficial impact on the new infection risk, especially when a foremilk cup was used. Keeping fresh-lactating cows in pens separated from the herd had a negative impact on the udder health, especially when housed together with sick cows. The results of this study confirm that the management of the milking routine and the environment can contribute to the control of udder health and milk quality.

**Key-words:** bovine mastitis, new infection risk, husbandry management

## Introduction

The health and productive performance of dairy cows is still significantly affected by the occurrence of inflammation of the udder because of intramammary infection with mastitis pathogens [1, 2]. Modern

milk quality programs focus on the prevention of new infections of the mammary gland to provide and maintain satisfactory and sustainable udder health [3, 4]. In Germany and other European countries, the new infection risk of lactating cows is used as a tool to estimate the level of new infections of the udder occurring at herd-level. This parameter is available monthly from dairy herd improvement tests (DHIT). The average new infection risk in dairy herds in lactation located in Lower Saxony, Northern German, the focus of this study, was 20.8% using a cutoff of 100,000 somatic cell counts per mL milk [5].

Prevention can only be sufficient if risk factors that can cause mastitis are identified and research has focused on this factor over the last decades [4, 6]. A lack of information about farm-level risk factors affecting the udder health of dairy herds in Germany has been published. In a large field study recently conducted in Northern Germany, environmental pathogens were found to be the major cause of clinical mastitis with *Streptococcus uberis* being the most prevalent pathogen [7]. Full cure rates were shown to be low, emphasizing the great relevance of preventing new infection and therefore also clinical mastitis [7]. Bedding material and bovine feces are considered as the main reservoirs and vectors of environmental pathogens causing intramammary infection. Different bedding materials were shown to be a source of exposure to environmental pathogens like *Klebsiella* spp. [8, 9] or *Streptococcus uberis* [8], but they can also be prevalent in feces [10, 11] as well as the indoor and outdoor housing environment [11, 12, 13]. Another recent German scientific paper concluded that bacterial exposure could be reduced by daily replacement of bedding material and frequent cleaning of the lying area, as well as pre-cleaning teats before milking and

dipping teats after milking [14].

Possible risk factors and thus also preventive measures are diverse [6], so that it is necessary to focus on the most important factors under the given farm-level conditions. The aim of this study in the context of a joint research project was to determine factors of husbandry management, which are associated with the new infection risk measured by DHIT in Northern German dairy herds.

## Material and Methods

### Herd Selection:

As part of the joint research project "SAM, Analysis of Dairy Production", 60 dairy farms were selected. According to the selection criteria of the joint research project, all of these had to be commercial farms located in Lower Saxony, Germany. The herds had to have at least 60 dairy cows of mainly Holstein breed kept in free stalls with cubicles and to take part in dairy herd improvement testing monthly to be eligible to participate. At the outset of this study, the number of cows housed on the farms ranged from 62 to 620 cows (arithmetic mean: 143.4; median: 116.5) and from 7,500 to 11,750 liters of milk yield per cow per year (arithmetic mean: 9,437.4; median: 9,500.0). The original topic of the joint research project was to compare indoor husbandry with pasture-based husbandry to identify differences in animal health and welfare, and specifically in udder health, while determining the factors of pasture management and husbandry that are associated with udder health parameters. Factors regarding pasture management as part of this research were analyzed separately and published in a previous paper [15]. Further details on material and methods are presented in that publication.

### Udder Health Data:

The new infection risk of lactating cows is defined for dairy herd improvement tests in Germany [16] as the percentage of lactating animals with > 100,000 somatic cells per mL milk of all lactating animals at monthly dairy herd improvement test having had ≤ 100,000 somatic cells per mL milk at the previous dairy herd improvement test. This parameter was calculated at farm-level using the individual somatic cell counts from monthly DHIT from January 2012 to April 2015. Arithmetic means of the new infection risk were determined at herd-level for the years 2012, 2013, 2014, the summer seasons (May to October) in 2012, 2013, 2014 and the winter seasons (November to April) in 2012/2013, 2013/14 and 2014/2015 based on the data provided monthly.

### Farm Data:

Each farm was visited by trained scientists at the beginning of the joint research project. On this occasion farm-specific data was documented following a standardized, structured questionnaire. The data regarding management of the dairy farm and its cattle was collected via interview of the farmer and investigating the livestock facilities (Table 1).

### Statistical Analysis:

For analyzing the dataset, the program SPSS 26.0, SPSS Inc. (Chicago, IL, USA) was used with the herd as the statistical unit. The subject was the herd (random) with repeated measurements. We found the unstructured covariance matrix structure to provide the best fit to these data. Associations between new infection risk (NIR) at herd-level level for the years 2012, 2013, 2014, the summer seasons (May to October) and the winter seasons (November to April) in 2012, 2013, 2014 and 2012/2013, 2013/14 and 2014/2015 (target) and risk factors (independent variables) were examined with generalized linear mixed models after pre-screening for variable selection in univariable analysis. The normal distribution of the outcome variable NIR was tested and confirmed using the Kolmogorov-Smirnow test. The relationship between

**Table 1: Farm-level variables considered in the univariable analysis of their associations with the new infection risk of lactating cows\*. Farm-level variables from the univariable analysis exhibiting a significant association ( $p \leq 0.1$ ) were considered in the multivariable analysis of their associations with the new infection risk of lactating cows\***

Category	Variable	$p \leq 0.1^{**}$
<b>Farm</b>		
performance	number of dairy cows	
	number of workers	**
	number of workers per dairy cow	
	average milk yield	
	length of productive life per cow	**
	average lifetime milk production per cow	**
	milk yield per length of productive life per cow	
	milk yield per life days per cow	
	replacement rate	
	age at first calving	**
calving interval		
management	regular pregnancy check	
	purchase or rearing of heifers	
	age of the farmer	
	years of job experience of the farmer	
	member of an advisory council	**
	use of dairy herd management software	
	udder health of cows is a priority	
	fertility of cows is a priority	
	hoof health of cows is a priority	
	longevity of cows is a priority	**
good overall health of cows is a priority		
milk yield per cow is a priority	**	
feed	forages are tested	
	grass silage is tested	
	corn silage is tested	
hoof health	prevalence of lameness diseases	
	prevalence of dermatitis digitalis	**
	prevalence of sole ulcers	**
	prevalence of white line disease	**
	prevalence of dermatitis interdigitalis and laminitis	**
	prevalence of dermatitis digitalis and white line disease	**
	number of hoof trimmings per cow per year	**
	hoof trimming is done by	
	method of hoof trimming	
	use of footbath	

udder health	most prevalent mastitis pathogen		hygiene	number of cows milked per milking cluster	
	number of milk samples for microbiological testing			milking-order of cow groups	
	species of mastitis pathogens prevalent in milk samples			milking-order of cows with mastitis	
				milking-order of fresh cows	
<b>Milking</b>				use of gloves in the milking parlor	**
technology	design of the milking parlor	**		cleaning of hands in the milking parlor	
	age of the milking parlor			sanitation of hands in the milking parlor	**
	maintenance interval of the milking machine	**		cleaning of teats before milking	**
	number of milking clusters	**		method of cleaning of teats before milking	**
	use of automatic cluster removers			number of cows per pre-cleaning-towel	**
	switch level of automatic cluster removers	**		cleaning of pre-cleaning towels	**
	assisted positioning of milking clusters			pre-milking teat sanitation	
	use of automatic post-milking stripping			post-milking teat sanitation	**
	milk yield recording			method of post-milking teat sanitation	**
	operating vacuum of milking machine	**		intermediate rinsing of clusters	
	type of pulsator			intermediate disinfection of clusters	**
	pulse ratio			removal of udder-hair	**
	design of the milk line	**		water quality in the milking parlor	**
	diameter of the milk line			cleaning of the milking stall floor	**
	diameter of the short milk hose			tools for cleaning the milking stall floor	**
	material of the short milk hose	**		cleansing of the milking machine	
	diameter of the long milk hose			temperature of rinsing water	**
	length of the long milk hose			alkaline cleansing of the milking machine per day	**
	material of the long milk hose			acid cleansing of the milking machine per day	**
	material of the liners				
	design of the liners	**			
	replacement interval of the liners followed according to specifications	**			
	sight glass in the milking parlor	**			
	air ingress during milking	**			
	driver in the waiting area				
work	who is milking				
	number of milkers per milking	**			
	parlor work routine	**			
	use of forestripping during pre-milking routine	**			
	examination of the milk during forestripping	**			
	preparation lag time	**			
	use of automatic stimulation				
	work-quality of attaching the milk clusters	**			
	use of oxytocin	**			

the NIR and the independent variables was first determined using appropriate univariable parametric test procedures. Independent variables associated with the dependent variable at  $p \leq 0.1$  except for predictors in the same model, which indicated a correlation of  $r > 0.70$  with one another (to avoid multicollinearity; for this reason, no variables were excluded) were submitted to generalized linear mixed models with an identity link.

The multivariable analysis was performed using a backward stepwise selection and elimination procedure until each independent variable

had a p-value of  $\leq 0.05$ . Confounding effects were monitored by observing regression coefficient changes. Variables that modified regression coefficients by  $> 20\%$  were considered confounding factors. No confounding was observed. The models were evaluated using the Akaike information criterion (AIC), where an AIC closest to zero was used as final model. In the final model, all biologically credible two-way interactions were tested but eliminated again due to lack of significance. Model fit was evaluated by checking normality of the residuals. The random farm effect was not significant in the models but was kept as a design variable. Least square means from the model were calculated. The significance level for the linear mixed model was 0.05.

## Results

### New Infection Risk:

Between January 2012 and April 2015, the monthly herd-level new infection risk of lactating cows varied between 7.4% and 43.0%. The arithmetic mean was 20.6% and the median 18.9%.

The monthly herd-level new infection risk of lactating cows calculated from individual somatic cell counts from monthly DHIT are shown in Table 2 for the respective years, summer (May to October) and winter seasons (November to April) in the period from January 2012 to April 2015.

### Univariable analysis:

Numerous factors of the management of the farm, milking, the lactating cows, the dry cows as well as the young stock were found to be associated with the new infection risk in the univariable analysis. In Table 1, the herd-specific factors that were statistically associated with the risk of new infection ( $p \leq 0.1$ ) in the univariable analysis are listed.

<b>Lactating cows</b>			<b>Dry cows</b>		
housing	number of groups of lactating cows	**	drying off	drying-off procedure	
	design of the barn	**		average length of the dry period	
	number of cows per stall			dry cow treatment	**
	number of cows per feed bunk space	**		method of dry cow treatment	
	housing area per cow			blanket dry cow treatment	
	design of the stalls	**	housing	number of groups of dry cows	**
	type of mattress in the stalls			design of the dry cow pen	**
	primary bedding material			design of the dry cow stalls	**
	secondary bedding material	**		type of mattress in the dry cow stalls	
	type of limestone added to the bedding material			bedding material in the dry cow stalls	**
	proportion of limestone of the bedding material			type of limestone added to the bedding material of dry cows	
	frequency of adding fresh bedding material			design of the housing floor in the dry cow pen	**
	frequency of cleaning the stalls			frequency of manure removal on housing floor of the dry cows	
	complete replacement of the bedding	**		method of manure removal on housing floor of dry cows	
	lactating cows on deep straw			access to pasture of dry cows	
	material of the housing floors	**		days per year access to pasture of dry cows	
	design of the housing floors	**		start of the pasture-season of the dry cows	**
	frequency of manure removal from the housing floor			end of the pasture-season of the dry cows	**
	method of manure removal from the housing floor	**		requirements (weather, grass growth) for access to pasture of dry cows	
	additional housing yards			design of the calving pens	
	access to pasture			number of calving pens	
	days per year access to pasture			average size of calving pens	
	hours per day access to pasture in March	**		overall size of the calving area	
	hours per day access to pasture in April			average square meters of calving area per cow	
	hours per day access to pasture in May			bedding material in the calving pens	**
	hours per day access to pasture in June			calving pens separated from sick cow pens	
hours per day access to pasture in July					
hours per day access to pasture in August	**		feed	number of dry cow rations	
hours per day access to pasture in September				separate ration fed to dry cows	
hours per day access to pasture in October					
hours per day access to pasture in November			<b>Young stock</b>		
access to pasture from November to April			calves	design of the calf pen in the first two weeks after birth	
selection gates				design of the calf pen after the first two weeks after birth	
consolidation of livestock trails				bedding material of the calves	
consolidation material of livestock trails				access to pasture of calves	
separate pen for fresh cows	**			age at first access to pasture of calves	
time in separate pen for fresh cows	**		young heifers	design of the young heifer pen	
				design of the young heifer stalls	
feed	source of drinking water			type of mattress in the young heifer stalls	
	access to additional feed of lactating cows if kept on pasture	**		bedding material of the young heifers	
	type of additional feed of lactating cows if kept on pasture			access to pasture of young heifers	
	type of watering place on pasture			days per year access to pasture of young heifers	
	number of different rations for lactating cows	**		age at first access to pasture of young heifers	

heifers	design of the heifer pen	
	design of the heifer stalls	
	type of mattress in the heifer stalls	
	bedding material of the heifers	**
	access to pasture of heifers	
	days per year access to pasture of heifers	
	age at first access to pasture of heifers	**
	age group at first access to pasture	**

\* percentage of lactating animals with > 100,000 somatic cells per mL milk of all lactating animals at monthly dairy herd improvement test having had ≤ 100,000 somatic cells per mL milk at the previous dairy herd improvement test

\*\* the respective variable was statistically associated ( $p \leq 0.1$ ) in the univariable analysis

#### Multivariable analysis:

Four different risk factors remained in the final generalized mixed models. These factors were related to milking and the environment of the lactating cows. The new infection risk was lower in herds with a pre-milking procedure including forestripping of every cow as a standard, and even lower if foremilk cups were used for this purpose compared to herds with no forestripping included in the pre-milking routine. If only animals which were conspicuous regarding udder diseases were forestripped, the new infection risk was higher than in those herds with no forestripping procedure. Very good quality work in attaching the milk-clusters to the udder was significantly associated with a lower new infection risk. The existence of fresh cow pens resulted in a higher new infection risk, especially if the fresh cows were housed together with the sick cows. The method of manure removal from housing floors provided significant results. Robots and vehicles with an attached scraper and manual scraping were associated with a lower new infection risk and manually controlled manure scrapers with a higher new infection risk compared to an automatic scraper. The results of the final multivariable analysis of the new infection risk of lactating cows including these risk factors are displayed in Table 3, and least square means describing differences between these risk factors in Table 4.

## Discussion

The aim of this study was to investigate the influence of manifold management factors on the new infection risk of dairy cattle in Lower Saxony, Germany. This was one of the first investigations focusing on such udder health risk factors in this geographic region. The regional impact on the results must be considered in the interpretation, even though no significant factors were region-specific in the narrower sense. It must also be recognized that the number of evaluated variables was high compared to the number of herds and respective data regarding udder health due to the given study design of the joint research project. Changes in the examined variables of the participating farms over the period of the investigation were not considered.

Despite this, the study provides evidence that management factors regarding milking and the environment of the dairy cows have an impact on udder health. Due to the limited power of the study, only the most important risk factors could probably be identified.

One of these factors is the pre-milking routine. Forestripping every cow before attaching the milking clusters had a beneficial effect on the new infection risk of the lactating cows, especially if the milk was collected with a foremilk cup. Forestripping only cows conspicuous of mastitis resulted in a higher risk of new infection.

Previous studies have shown a positive impact of using a foremilk cup on somatic cell counts at cow level [17] in the forestripping practice, but a negative impact on the number of cases of clinical mastitis [18] and clinical mastitis caused by *Staphylococcus aureus* [19, 20, 21] and *Escherichia coli* [20] when cows were forestripped manually. More recent investigations either found no association between forestripping during milking preparation and the somatic cell count of the bulk tank milk [22] or showed a tendency for forestripping to reduce the occurrence of clinical mastitis and a significant reduction thereof if forestripping was part of a complete milking routine (forestripping, predipping and drying before milking unit attachment) [23].

The procedure of forestripping could also be related to the preparation lag time, which was only a significant factor in the univariable analysis. Preparation lag time was associated with clinical and subclinical mastitis [24, 25].

Forestripping is a recommended practice for a complete milking routine as well as mastitis control programs [3, 23, 26] and is a mandatory practice by legislation of the European Union [27]. Nonetheless, the

**Table 2: Farm-level new infection risk of lactating cows\* [%] calculated from individual somatic cell counts from monthly DHIT in the period from January 2012 to April 2015**

Period	Minimum	Lower Quartile	Median	Upper Quartile	Maximum	Mean
All	7.4	15.1	18.9	25.2	43.0	20.6
2012	10.0	17.0	22.0	28.0	39.0	22.6
2013	8.0	15.0	18.0	24.0	35.0	19.7
2014	10.0	15.0	18.0	24.3	37.0	19.7
2012S	8.1	18.7	23.5	30.8	8.1	25.0
2013S	7.4	16.6	18.6	23.8	7.4	20.7
2014S	10.4	15.2	18.6	25.2	10.4	20.8
2012/2013W	8.8	13.1	17.1	21.5	8.8	18.2
2013/2014W	8.7	14.1	17.8	25.3	8.7	19.8
2014/2015W	8.0	14.0	17.3	23.7	8.0	18.9

\* percentage of lactating animals with > 100,000 somatic cells per mL milk of all lactating animals at monthly dairy herd improvement test having had ≤ 100,000 somatic cells per mL milk at the previous dairy herd improvement test

S = summer-season (May to October)

W = winter-season (November to April)

**Table 3: Final multivariable analysis of the new infection risk of lactating cows\***

Variable	Coefficient	Standard Error	Confidence interval 2,5%	Confidence interval 97,5%	p
<b>Intercept</b>	28.328	2.052	24.306	32.349	0.000
<b>Use of forestripping during pre-milking routine</b>					
Yes	-5.094	1.252	-7.548	-2.640	0.000
Yes, cup is used	-6.151	1.238	-8.576	-3.725	0.000
Yes, only conspicuous cows	13.179	2.373	8.527	17.830	0.000
No	0				
<b>Work-quality of attaching milk clusters</b>					
Very good	-5.412	1.270	-7.901	-2.922	0.000
Good	-1.046	1.300	-3.594	1.503	0.421
Satisfactory	2.931	2.289	-1.554	7.417	0.200
Just sufficient	0				
<b>Method of manure removal from housing floors</b>					
Manual scraping	-5.103	1.796	-8.622	-1.584	0.004
Robot	-6.650	2.221	-11.003	-2.297	0.003
Vehicle	-4.257	1.547	-7.289	-1.225	0.006
Manually controlled scraper	5.653	1.416	2.878	8.428	0.000
None	-2.307	1.908	-6.047	1.433	0.227
Automatic scraper	0				
<b>Separate pen for fresh cows</b>					
Yes	3.426	1.243	0.990	5.863	0.006
Yes, kept together with sick cows	5.350	1.558	2.297	8.404	0.001
No	0				

\*percentage of lactating animals with > 100,000 somatic cells per mL milk of all lactating animals at monthly dairy herd improvement test having had ≤ 100,000 somatic cells per mL milk at the previous dairy herd improvement test

quality of how it is performed in terms of pre-milking hygiene, especially when contagious mastitis pathogens are prevalent, could be crucial for the outcome.

Another finding associated with the milking practice was a lower new infection risk related to high-quality work in attaching the milk-clusters to the udder. Quality indicators included the extent of air admission, hygiene and the speed of attaching the milk-clusters. A French study [28] showed a negative impact of air admission at teat cup attachment, this being an indication of low-quality work. Air admission could lead to

a varying vacuum in the milking system and even fall-offs of the milking clusters with poorer udder health resulting [29].

The method of manure removal from the housing floors of the lactating cows was statistically significant in the final model of our study. The methods were recorded as manual scraping, robot, vehicle, manually controlled scraper, none and automatic scraper, with the manually controlled scraper having the highest risk of new infection in lactation and the robot having the lowest risk. Possibly the method of manure removal refers to the new infection risk if it has a sufficient impact on

**Table 4: Least square means describing differences between farm-level variables associated with the new infection risk of lactating cows\***

Variable	Mean	Standard Error	Confidence interval 2,5%	Confidence interval 97,5%
<b>Use of forestripping during pre-milking routine</b>				
Yes	18.527	0.737	17.081	19.970
Yes, cup is used	17.470	0.990	15.529	19.410
Yes, only conspicuous cows	36.799	1.802	33.266	40.332
No	23.620	1.193	21.281	25.960
<b>Work-quality of attaching milk clusters</b>				
Very good	19.573	0.932	17.747	21.400
Good	23.939	0.641	22.683	25.196
Satisfactory	27.916	1.815	24.359	31.474
Just sufficient	24.985	1.287	22.463	27.507
<b>Method of manure removal from housing floors</b>				
Manual scraping	21.111	1.554	18.065	24.157
Robot	19.564	1.443	16.737	22.393
Vehicle	21.957	1.564	19.690	24.223
Manually controlled scraper	31.867	1.288	29.342	34.393
Automatic scraper	26.214	1.199	23.864	28.565
None	23.907	1.651	20.670	27.144
<b>Separate pen for fresh cows</b>				
Yes	29.245	1.180	26.933	31.558
Yes, kept together with sick cows	31.169	1.407	28.412	33.927
No	25.819	0.629	24.587	27.051

\* percentage of lactating animals with > 100,000 somatic cells per mL milk of all lactating animals at monthly dairy herd improvement test having had ≤ 100,000 somatic cells per mL milk at the previous dairy herd improvement test

the hygiene of the environment. Numerous other studies could show a relation between the hygiene of floors and passages as well as udder health indicators [30, 31, 32, 33].

The collected data provided evidence that the existence of a pen for fresh cows was statistically associated with a higher new infection risk. No references with the same or with differing findings are available. The study dataset did not provide clear information about the conditions and the management of the fresh cow pens. However, the dataset provided information that it was a common practice to house fresh cows in group pens on deep straw during the first days after calving in at least 29 of the 60 farms. This type of housing could be associated with poorer udder health because of worse environmental hygiene compared to free stall housing [18, 34, 35].

Several investigations stated the importance of optimal environmental hygiene for the transition area [31] and the calving area [18, 19, 20, 21, 30, 31, 36, 37]. The environmental hygiene fresh cows are exposed to after calving could be of similar importance.

When the fresh cows were housed together with the sick cows, the described effect was intensified. This practice possibly increases the exposure to mastitis pathogens. Comparable results were stated concerning the incidence rate of clinical mastitis caused by *Escherichia coli* when calving pens and sick cow pens were not separated [19].

## Conclusions

Farm-level risk factors for new infections of the udder of dairy cows are multitudinous and depend on the given conditions of the respective farm. Minimizing the new infection risks can be effectively achieved by optimizing the management of the milking routine and the environment that cows are housed in. The results of this research suggest keeping fresh cows under hygienic conditions and separated from sick cows and to include forestripping to the pre-milking routine. The use of a foremilk cup can be a purposeful tool to maximize hygiene in this practice.

## Acknowledgment

The study was supported by the Ministry for Science and Culture of Lower Saxony (MWK) within the collaborative research project SAM, Analysis of Dairy Production: Grazing versus Indoor Housing of Dairy Cows, Support Code: ZN 2864.

## References

- Halasa T, Huijps K, Østeras O, Hogeveen H. Economic effects of bovine mastitis and mastitis management: A review. *Veterinary Quarterly*, 2007; 29(1): 18–31.
- Seegers H, Fourichon C, Beaudeau F. Production effects related to mastitis and mastitis economics in dairy cattle herds. *Vet Res*, 2003; 34: 475–491.
- Ruegg PL. Managing cows, milking and the environment to minimize mastitis. *Japanese Journal of Large Animal Clinics*, 2015; 5(4):210.
- Ruegg PL. A 100-Year Review: Mastitis detection, management, and prevention. *J Dairy Sci*, 2017; 100:10381–10397.
- Behr B, Hachenberg S. Strategisches Eutergesundheitsmonitoring – erste Ergebnisse einer deutschlandweiten Umsetzung. *Proc. BPT Kongress 2016*.
- Volling O, Krömker V. Udder health management practices in dairy enterprises to reduce the incidence of bovine mastitis. *Dtsch Tierärztl Wochenschr*, 2008; 115(11): 410–420.
- Schmenger A, Krömker V. Characterization, cure rates and associated risks of clinical mastitis in Northern Germany *Vet Sci*, 2020; 7: 170.
- Ericsson Unnerstad H, Lindberg A, Persson Waller K, Ekman T, Artursson K, Nilsson-Öst M, Bengtsson B. Microbial aetiology of acute clinical mastitis and agent-specific risk factors. *Vet Microbiol*, 2009; 137: 90–97.
- Munoz MA, Welcome FL, Schukken YH, Zadoks RN. Molecular epidemiology of two *Klebsiella pneumoniae* mastitis outbreaks on a dairy farm in New York State. *Journal of Clinical Microbiology*, 2007; 45(12): 3964–3971.
- Munoz MA, Ahlström C, Rauch BJ, Zadoks RN. Fecal shedding of *Klebsiella pneumoniae* by dairy cows. *J Dairy Sci*, 2006; 89(9): 3425–3430.
- Zadoks R, Tikofsky LL, Boor KJ. Ribotyping of *Streptococcus uberis* from a dairy's environment, bovine feces and milk. *Veterinary Microbiology*, 2005; 109(3–4): 257–265.
- Lopez-Benavides MG, Williamson JH, Pullinger GD, Lacy-Hulbert SJ, Cursons RT, Leigh JA. Field observations on the variation of *Streptococcus uberis* populations in a pasture-based dairy farm. *J Dairy Sci*, 2007; 90: 5558–5566.
- Olde Riekerink RGM, Barkema HW, Stryhn H. The effect of season on somatic cell count and the incidence of clinical mastitis. *J Dairy Sci*, 2007; 90: 1704–1715.
- Hohmann MF, Wente N, Zhang Y, Krömker V. Bacterial load of the teat apex skin and associated factors at herd level. *Animals*, 2020; 10: 1647.
- Gösling M, Klocke D, Reinecke F, Zoche-Golob V, tho Seeth M, Paduch JH, Krömker V. Pasture-associated influence on the udder health of dairy herds in Northern Germany. *Milk Sci Int*, 2019; 72: 2–10.
- DLQ (German Association for Performance and Quality Testing). DLQ-Richtlinie 1.15: Zur Definition und Berechnung von Kennzahlen zum Eutergesundheitsmonitoring in der Herde und von deren Vergleichswerten, 2014; DLQ guideline 1.15, Bonn, Germany.
- Gill R, Howard WH, Leslie KE, Lissemore K. Economics of mastitis control. *J Dairy Sci*, 1990; 73: 3340–3348.
- Peeler EJ, Green MJ, Fitzpatrick JL, Morgan KL, Green LE. Risk factors associated with clinical mastitis in low somatic cell count British dairy herds. 9th International Symposium on Veterinary Epidemiology and Economics, 2000.
- Barkema HW, Schukken YH, Lam TJ, Beiboer ML, Benedictus G, Brand A. Management practices associated with the incidence rate of clinical mastitis. *J Dairy Sci*, 1999; 82: 1643–1654.
- Elbers ARW, Miltenburg JD, De Lange D, Crauwels APP, Barkema HW, Schukken YH. Risk factors for clinical mastitis in a random sample of dairy herds from the southern part of the Netherlands. *J Dairy Sci*, 1998; 81: 420–426.
- Schukken YH, Grommers FJ, Van de Geer D, Ere HN, Brand A. Risk factors for clinical mastitis in herds with a low bulk milk somatic cell count. 2. Risk factors for *Escherichia coli* and *Staphylococcus aureus*. *J Dairy Sci*, 1991; 74: 826–832.
- Belage E, Dufour S, Bauman C, Jones-Bitton A, Kelton DF. The Canadian National Dairy Study 2015 — Adoption of milking practices in Canadian dairy herds. *J Dairy Sci*, 2017; 100: 3839–3849.
- Rodrigues ACO, Caraviello DZ, Ruegg PL. Management of Wisconsin dairy herd enrolled in milk quality teams. *J Dairy Sci*, 2005; 88: 2660–2671.
- Giovannini G, Zecconi A. Field study on epidemiology of clinical

- mastitis in five Italian dairy herds. *Milk Sci Int*, 2002; 57: 3–6.
25. Kawai K, Kurosawa S, Nagahata H, Rosenberg J. Evaluation of dairy management practices on Japanese dairy farms: relationships to bulk tank SCC and individual linear scores. *NMC Annual Meeting*, 2005; 285–286.
  26. Pankey JW, Wildman EE, Drechsler PA, Hogan JS. Field trial evaluation of premilking teat disinfection. *J Dairy Sci*, 1987; 70: 867–872.
  27. European Commission. Regulation (EC) No 853/2004 of the European Parliament and of the Council of 29 April 2004 laying down specific hygiene rules for food of animal origin. *Official Journal of the European Communities*, 2004; L139/55.
  28. Chassagne M, Barnouin J, Le Guenic M. Expert assessment study of milking and hygiene practices characterizing very low somatic cell score herds in France. *J Dairy Sci*, 2005; 88: 1909–1916.
  29. Tan J, Janni KA, Appleman RD1993. Milking system dynamics. 2. Analysis of vacuum systems. *J Dairy Sci*, 1993; 76: 2204–2212.
  30. Barnouin J, Chassagne M, Bazin S, Boichard D. Management practices from questionnaire surveys in herds with very low somatic cell score through a national mastitis program in France. *J Dairy Sci*, 2004; 87: 3989–3999.
  31. Green MJ, Bradley AJ, Medley GF, Browne WJ. Cow, farm, and management factors during the dry period that determine the rate of clinical mastitis after calving. *J Dairy Sci*, 2007; 90: 3764–3776.
  32. Kelly PT, O’Sullivan K, Berry DP, More SJ, Meaney WJ, O’Callaghan EJ, O’Brien B. Farm management factors associated with bulk tank somatic cell count in Irish dairy herds. *Ir Vet J*, 2009; 62: 45–51.
  33. Volling O, Krömker V, Brinkmann J, March S, Weiler M, Weiß M, Drerup C, Becker M, Klocke D, Merch F. Haltungshygiene und Eutergesundheit in ökologisch geführten Milchviehbetrieben. 11. Wissenschaftstagung Ökologischer Landbau, 2011; Tagungsband, Band 2.
  34. Barnouin J, Bord S, Bazin S, Chassagne M. Dairy management practices associated with incidence rate of clinical mastitis in low somatic cell score herds in France. *J Dairy Sci*, 2005; 88: 3700–3709.
  35. Richert RM, Cicconi KM, Gamroth MJ, Schukken YH, Stiglbauer KE, Ruegg PL. Risk factors for clinical mastitis, ketosis, and pneumonia in dairy cattle on organic and small conventional farms in the United States. *J Dairy Sci*, 2013; 96: 4269–4285.
  36. Green MJ, Bradley AJ, Medley GF, Browne WJ. Cow, farm, and herd management factors in the dry period associated with raised somatic cell counts in early lactation. *J Dairy Sci*, 2008; 91: 1403–1415.
  37. Nyman AK, Emanuelson U, Gustafsson AH, Persson Waller K. Management practices associated with udder health of first-parity dairy cows in early lactation. *Prev Vet Med*, 2009; 88(2): 138–149.

*Copyright © 2021 Milk Science International. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY) 4.0. The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.*